Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the joint a very interesting and peculiar mechanical structure. There are no ligamenta alaria or mucosa.

May 28.

Mr. Thomas Meehan, Vice-President, in the chair.

Twenty-three persons present.

On Polyxenes fasciculatus.—Mr. J. A. **Ryder** announced that he had identified the myriapod *Polyxenes fasciculatus*, Say, in the vicinity of the city at several places in the park. It was found that its morphological history agreed with that of *P. lagurus*, of Europe, as detailed by Bode,[1] but there was the same paucity of males as observed by that naturalist. The presence of the hooks terminating the caudal bristles was an independent discovery so far as the American species was concerned, Mr. Ryder not having seen Bode's paper until three days after his own observations had been made. Upon comparing these parts of the two species together we find the ends of the bristles, according to Bode, in the European species bent in the form of a semicircle, with the subterminal processes depending parallel to each other from within the arc; in our species, on the other hand, the bristle terminates in two barbed divaricating points directed forwards, forming a subacute angle with the supporting shaft of the bristle, the subterminal processes depending within the angle on the supporting shaft next the animal. This new locality indicates a wide distribution for the animal, viz., from Georgia, where it was first discovered by Say, to Massachusetts, where it has been found by Packard.

Corundum in North Carolina.—**Joseph Willcox** said that he desired to place on record a locality for Corundum recently discovered in Iredell Co., North Carolina. Near Statesville, as the southern limit, it is found associated with Kyanite and Serpentine. It is also found at several localities for a distance of twenty miles nearly northeast of Statesville, and at its northern limit it is associated with Steatite, Actinolite, and Damourite. No large specimens of Corundum have yet been found at this locality.

Prof. Gumesindo Mendoza, of Mexico, and Stephen Bowers of Santa Barbara, Cal., were elected Correspondents.

The following paper was ordered to be printed:—

---

[1] Zeitsch. f. die gesammten Naturwissenschaften. 3te folge, Berlin, 1877. Bd. II., 8vo. pp. 233-268, Pl. 11-14.